

Randomisation Program

Call Freefone 0800 138 5451 to contact the RECOVERY team for URGENT problems using the Randomisation Program or for medical advice. All NON-URGENT queries should be emailed to recoverytrial@ndph.ox.ac.uk

Logged in as: RECOVERY Site

Section A: Baseline and Eligibility

Date and time of randomisation: 27 Mar 2022 14:00

Treating clinician
A1. Name of treating clinician

Patient details
A2. Patient surname
 Patient forename

A3. NHS number Tick if not available

A4. What is the patient's date of birth? / / Age: 22y 2m

A5. What is the patient's sex?

Inclusion criteria
A6. Has consent been taken in line with the protocol? Tick if not available
If answer is no patient cannot be enrolled in the study
 NB current PIS/ICF version is V22.0 (adults) or V14.0 (children)

A6.0.1 How was consent obtained?

A6.5 Does this patient have viral pneumonia? Yes

See protocol for typical features. If answer is no patient cannot be enrolled in the study

A7.0 Does the patient have proven SARS-CoV-2 infection? Yes

A7.0.1 What was lateral flow test result?

A7.0.2 What was PCR test result?

A7.1 Does the patient have proven influenza infection?

A8. Does the patient have any medical history that might, in the opinion of the attending clinician, put the patient at significant risk if they were to participate in the trial?

A9. Symptom onset date: / /

A10. Date of hospitalisation: / /

A11. Does the patient require oxygen?

A12. Please select one of the following to describe the current level of ventilation support

A12.1 Enter latest oxygen saturation measurement (%)

A12.2 Enter latest CRP measurement since admission to hospital (mg/L) Tick if not measured
Enter 0 if below the limit of measurement Tick if greater than limit of measurement

A12.3 Enter latest creatinine measurement since admission to hospital Tick if not measured
Enter 0 if below the limit of measurement Tick if greater than limit of measurement

A12.4 Enter latest D-dimer measurement since admission to hospital Tick if not measured
Enter 0 if below the limit of measurement Tick if greater than limit of measurement

A12.5 Has the patient received a COVID-19 vaccine?

A12.6 Has the patient received an influenza vaccine in the last 12 months?

Does the patient have any CURRENT comorbidities or other medical problems or treatments?

A13.1 Diabetes

A13.2 Heart disease

A13.3 Chronic lung disease

A13.4 Tuberculosis

A13.5 HIV

A13.6 Severe liver disease

A13.7 Severe kidney impairment (eGFR < 30 or on dialysis)

A13.9.0 Does their clinician consider the patient to be severely immunocompromised?

A13.12 Has the patient received tocilizumab or sarilumab therapy during this admission?

A13.14 Current or planned treatment with neuraminidase inhibitor
eg, oseltamivir, zanamivir

A13.15 Has the patient received casirivimab+imdevimab (Ronapev) during this illness?

A13.16 Has the patient received sotrovimab during this illness?

A13.17 Has the patient received molnupiravir during this illness?

A13.18 Has the patient received Paxlovid during this illness?

Are the following treatments UNSUITABLE for the patient? If you answer Yes, it means you think this patient should NOT receive this drug.

A14E.1 High-dose corticosteroids Tick if not suitable
NB Please carefully consider suitability of patients already on higher doses (>= 7mg/day dexamethasone or equivalent). Patients eligible for the Paxlovid comparison will be automatically marked as unsuitable for this comparison.

A14F.1 Empagliflozin Tick if not suitable
Empagliflozin is NOT suitable if patient (i) has type 1 or post-pancreatotomy diabetes mellitus; or (ii) has a history of ketoacidosis; or (iii) has blood ketones >= 1.5 mmol/L or urine ketones 2+; or (iv) is pregnant or breastfeeding. Empagliflozin cannot be given via an enteral feeding tube.

A14J.1 Sotrovimab

A14K.1 Molnupiravir Tick if not suitable
NB Molnupiravir is NOT suitable if patient cannot swallow capsules.

A14L.1 Paxlovid No Tick if not suitable
NB Paxlovid contains ritonavir and has many drug-drug interactions (see protocol and SmPC). Please ensure these have been checked. Paxlovid is NOT suitable if patient cannot swallow tablets. Paxlovid is not suitable for pregnant women in the first trimester.

Are the following treatments available?

A15E.1 High-dose corticosteroids

A15F.1 Empagliflozin

A15J.1 Sotrovimab

A15K.1 Molnupiravir

A15L.1 Paxlovid Yes

Current medication
A16.1 Is the patient currently prescribed remdesivir?

A16.2 Is the patient currently prescribed systemic corticosteroids (dexamethasone, prednisolone, hydrocortisone, methylprednisolone)? Tick if not available
Please do not include topical or inhaled treatments

A16.5 What venous thromboembolism prophylaxis is the patient receiving? Tick if not available
Standard = usual for hospitalised patients (not increased due to COVID-19); Higher dose = treatment dose or increased prophylaxis due to COVID-19, or oral anticoagulation (eg, warfarin/DOAC).

A16.6 Is the patient currently prescribed baricitinib (or other JAK inhibitor)?

Serum sample collection
A17.0 Please confirm that patient has had a baseline serum sample collected according to the protocol

A17.1 Please confirm that patient has had a baseline nasal swab collected according to the protocol

Please sign off this form once complete
 Surname:
 Forename:
 Professional email: