

Randomisation Program

Call Freephone 0800 138 5451 to contact the RECOVERY team for URGENT problems using the Randomisation Program or for medical advice. All NON-URGENT queries should be emailed to recoverytrial@ndph.ox.ac.uk

Logged in as: RECOVERY Site

Section A: Baseline and Eligibility

Date and time of randomisation: 28 Mar 2022 14:00

Treating clinician
A1. Name of treating clinician

Patient details
A2. Patient surname
 Patient forename
A3. National ID Tick if not available

A4. What is the patient's date of birth? 01 / January / 2000 Age: 22y 2m
A5. What is the patient's sex?

Inclusion criteria
A6. Has consent been taken in line with the protocol?
If answer is No patient cannot be enrolled in the study
A6.0.1 How was consent obtained?
A6.5 Does this patient have viral pneumonia?
See protocol for typical features. If answer is No patient cannot be enrolled in the study
A7.0 Does the patient have proven SARS-CoV-2 infection?
A7.0.1 What was lateral flow test result?
A7.0.2 What was PCR test result?
A8. Does the patient have any medical history that might, in the opinion of the attending clinician, put the patient at significant risk if they were to participate in the trial?
A9. Symptom onset date: / /
A10. Date of hospitalisation: / /
A11. Does the patient require oxygen?
A12. Please select one of the following to describe the current level of ventilation support
A12.1 Enter latest oxygen saturation measurement (%)
A12.2 Enter latest CRP measurement since admission to hospital (mg/L) Tick if not measured
Enter 0 if below the limit of measurement Tick if greater than limit of measurement
A12.3 Enter latest creatinine measurement since admission to hospital mg/dL Tick if not measured
Please select correct units for the measurement (mg/dL or µmol/L)
A12.4 Enter latest D-dimer measurement since admission to hospital mg/L Tick if not measured
Please select correct units for the measurement (mg/L or ng/mL). Enter 0 if below the limit of measurement Tick if greater than limit of measurement
A12.5 Has the patient received a COVID-19 vaccine?
A12.6 Has the patient received an influenza vaccine in the last 12 months?
Does the patient have any CURRENT comorbidities or other medical problems or treatments?
A13.1 Diabetes
A13.2 Heart disease
A13.3 Chronic lung disease
A13.4 Tuberculosis
A13.5 HIV
A13.6 Severe liver disease
A13.7 Severe kidney impairment (eGFR<30 or on dialysis)
A13.7.1 Is the patient on dialysis or haemofiltration?
A13.9.0 Does their clinician consider the patient to be severely immunocompromised?
A13.12 Has the patient received tocilizumab or sarilumab therapy during this admission?
A13.14 Current or planned treatment with neuraminidase inhibitor
eg, oseltamivir, zanamivir
A13.15 Has the patient received casirivimab+imdevimab (Ronapreve) during this illness?
A13.16 Has the patient received sotrovimab during this illness?
A13.17 Has the patient received molnupiravir during this illness?
A13.18 Has the patient received Paxlovid during this illness?
Are the following treatments UNSUITABLE for the patient? If you answer Yes it means you think this patient should NOT receive this drug.
A14E.1 High-dose corticosteroids
A14F.1 Empagliflozin
Empagliflozin is NOT suitable if patient (i) has type 1 or post-pancreatectomy diabetes mellitus; or (ii) has a history of ketoacidosis; or (iii) has blood ketones ≥ 1.5 mmol/L or urine ketones ≥ 2+; or (iv) is pregnant or breastfeeding. Empagliflozin cannot be given via an enteral feeding tube.
A14K.1 Molnupiravir
Molnupiravir is NOT suitable if patient cannot swallow capsules.
Are the following treatments available?
A15E.1 High-dose corticosteroids
A15F.1 Empagliflozin
A15K.1 Molnupiravir
Current medication
A16.1 Is the patient currently prescribed remdesivir?
A16.2 Is the patient currently prescribed systemic corticosteroids (dexamethasone, prednisolone, hydrocortisone, methylprednisolone)?
Please do not include topical or inhaled treatments
A16.5 What venous thromboembolism prophylaxis is the patient receiving?
Standard = usual for hospitalised patients (not increased due to COVID-19); Higher dose = treatment dose or increased prophylaxis due to COVID-19, or oral anticoagulation (eg, warfarin/DOAC).
A16.6 Is the patient currently prescribed baricitinib (or other JAK inhibitor)?
Please sign off this form once complete
 Surname:
 Forename:
 Professional email: