

Clinical Area Preparation Record – Sotrovimab 1000mg in 100mL Sodium Chloride 0.9% Infusion Bag (Total volume = 116 mL)

Set up	Preparation of Infusion Bag				
<p>Step 1 Remove from the refrigerator:</p> <ul style="list-style-type: none"> 2 x Sotrovimab 500mg (62.5mg/mL) Concentrate for Solution for Infusion vial. <p>Select:</p> <ul style="list-style-type: none"> 1 x Sodium Chloride 0.9% 100mL Infusion Bag 1 x 20mL luer lock syringe 1 x drawing up needle 1 x 0.2 micron administration filter 	<p>Step 1 Bring the Sodium Chloride 0.9% 100mL Infusion Bag from the left side of the preparation area into the middle, swab the bung with a sterile 70% alcohol wipe and allow to dry.</p>				
<p>Step 2 Visually inspect the Sotrovimab vials</p> <ul style="list-style-type: none"> The solution should be clear, colourless or yellow to brown and free from visible particles <p>Should particulate matter or discoloration be observed, the vial must be discarded and replaced with a new vial.</p>	<p>Step 2 Bring the Sotrovimab 500mg vials from the left side of the preparation area into the middle, swab the bungs with sterile 70% alcohol wipe and allow to dry.</p>				
<p>Step 3 Place to the left side of the preparation area:</p> <ul style="list-style-type: none"> 2 x Sotrovimab 500mg vial 1 x Sodium Chloride 0.9% 100ml Infusion Bag 	<p>Step 3 Gently swirl the vials several times before use without creating air bubbles. Do not shake or vigorously agitate the vial.</p>				
<p>Step 4 Prepare an infusion additive label with the following details:</p> <ul style="list-style-type: none"> Sotrovimab 1000mg in Sodium Chloride 0.9% (Total volume = 116 mL) Date and time prepared [Additional details as required by local label format] 	<p>Step 4 Attach a drawing up needle to a 20mL luer lock syringe and draw up 1 x 16mL of Sotrovimab 500mg (62.5mg/mL) from the 2 vials</p> <p>Step 5 Add 16mL of Sotrovimab (62.5mg/mL) to the Sodium Chloride 0.9% 100mL Infusion Bag. Discard the syringe and needle into a yellow lidded sharps bin</p>				
	<p>Step 6 Gently rock the infusion bag back and forth 3 to 5 times. NB: Do not invert the infusion bag. Avoid forming air bubbles. Do not shake</p>				
	<p>Step 7 Attach the pre-prepared label to the bag</p>				
	<p>Step 8 Ensure the product is administered using an [Insert local in-line or add-on 0.2µm filter used] as a single IV infusion for 60 minutes</p>				
	<p>Step 9 Record details of the patient who will receive the bag below, and file the completed <i>Clinical Area Preparation Record</i> in accordance with local guidance</p>				
	Patient Name		Hospital No		Date of Birth

Document	RECOVERY - Clinical Area Preparation Record - Sotrovimab 1000mg in 100mL NaCl 0.9% Infusion Bag	Version Number	1	Date Issued	24/12/2021	Issued By	RECOVERY
		Site Name:		Review Date	14/12/2023	Approved by	