

Screening

PI or delegate identifies potential participant



Obtain ICF
(patient, witness or LAR)

Eligibility

- (i) Hospitalised (planned overnight stay) and aged ≥ 18 years old
- (ii) Pneumonia syndrome (*criteria below are a guide only - diagnosis is based on clinical opinion of the managing doctor*)
 - a) typical symptoms of a new respiratory infection; AND
 - b) objective evidence of acute lung disease (e.g. X-ray/CT, hypoxia, compatible clin. examination); AND
 - c) alternative causes are unlikely or excluded (e.g. heart failure)
- (iii) No medical history that might put patient at significant risk if participating in the trial
- (iv) Clinician does not believe a specific trial treatment is indicated or contra-indicated
- (v) One of the following diagnoses:

Confirmed influenza A or B

Patients can enter all three comparisons if eligible

Open to all patients weighing ≥ 40 kg except if pregnant or breastfeeding

Open to all patients

OR

CAP with planned antibiotic treatment

(**without** suspected SARS-CoV-2, influenza, active pulmonary tuberculosis or *Pneumocystis jirovecii* pneumonia)

Open for patients:

- 1) with hypoxia (on O₂ or Sats $<92\%$)
- 2) without suspected SARS-CoV-2

Baloxavir OR usual care *without* baloxavir

Oseltamivir OR usual care *without* neuraminidase inhibitor

Dexamethasone (6 mg daily)[†] OR usual care *without* systemic corticosteroids

Dexamethasone (6 mg daily)[†] OR usual care *without* systemic corticosteroids

[†]prednisolone/hydrocortisone in case of pregnancy/breastfeeding

Refer to Intervention Sheets for detailed treatment and dosing guide, available at www.recoverytrial.net/eu

Not all treatment comparisons may be open at your site – check with the study team



RECOVERY

Contact details local trial team

Name:

Email:

Phone number:

Name:

Email:

Phone number:

Name:

Email:

Phone number: