

RECOVERY

Randomised Evaluation of COVID-19 Therapy

Statistical Analysis Plan

Version 2.1

Date: 02 December 2020

Aligned with protocol version: 11.1, 21 November 2020

IRAS no: 281712

REC ref: EE/20/0101

ISRCTN: 50189673

EudraCT: 2020-001113-21

Nuffield Department of
POPULATION HEALTH



Table of Contents

Table of Contents.....	2
Abbreviations.....	5
List of authors and reviewers (up to and including SAP version 1.1)	6
List of authors and reviewers (version 2.0 onwards)	6
Roles and responsibilities	7
1 Introduction	8
2 Background information	8
2.1 Rationale	8
2.2 Objectives of the trial.....	9
2.2.1 Primary objective	9
2.2.2 Secondary objectives	9
2.3 Trial design	9
2.4 Eligibility	9
2.4.1 Inclusion criteria.....	9
2.4.2 Exclusion criteria	9
2.5 Treatments	9
2.5.1 Main randomisation part A:.....	9
2.5.2 Main randomisation part B:.....	10
2.5.3 Main randomisation part C:.....	10
2.5.4 Second randomisation for patients with progressive COVID-19.....	10
2.6 Definitions of primary and secondary outcomes.....	11
2.6.1 Primary outcome	11
2.6.2 Secondary clinical outcomes.....	11
2.6.3 Subsidiary clinical outcomes.....	11
2.6.4 Safety outcomes	11
2.6.5 Detailed derivation of outcomes	12
2.9 Randomisation	12
2.9.1 Main randomisation part A.....	12
2.9.2 Main randomisation part B.....	13
2.9.3 Main randomisation part C.....	13
2.9.4 Second randomisation for patients with progressive COVID-19.....	13
2.10 Blinding.....	14
2.11 Data collection schedule.....	14
2.12 Data monitoring.....	14

2.13	Trial reporting	14
3	Analysis populations	15
3.1	Population definitions	15
4	Descriptive analyses.....	15
4.1	Participant throughput.....	15
4.2	Baseline comparability of randomised groups	15
4.2.1	Main randomisation (parts A, B and C).....	15
4.2.2	Second randomisation	16
4.3	Completeness of follow-up	16
4.4	Adherence to treatment	16
5	Comparative analyses	16
5.1	Main randomisation part A	17
5.1.1	Primary outcome	17
5.1.2	Secondary outcomes.....	17
5.1.2.1	Time to discharge alive from hospital.....	17
5.1.2.2	Use of invasive mechanical ventilation (including ECMO) or death	17
5.1.3	Subsidiary clinical outcomes	17
5.1.3.1	Use of ventilation (overall and by type).....	17
5.1.3.2	Duration of invasive mechanical ventilation (time to successful cessation of invasive mechanical ventilation).....	18
5.1.3.3	Use of renal dialysis or haemofiltration	18
5.1.3.4	Thrombotic event.....	18
5.2	Main randomisation part B	18
5.3	Main randomisation part C	18
5.4	Second randomisation	18
5.5	Pre-specified subgroup analyses.....	19
5.6	Sensitivity analyses.....	19
5.7	Other exploratory analyses	19
5.8	Adjustment for baseline characteristics	20
5.9	Significance levels and adjustment of p-values for multiplicity.....	20
5.10	Statistical software employed	20
5.11	Data standards and coding terminology	20
6	Safety data	20
6.1.1.1	Cause-specific mortality	21
6.1.1.2	Major cardiac arrhythmia.....	21
6.1.1.3	Major bleeding	21

6.1.1.4	Early safety of antibody-based therapy	21
7	Additional POST-HOC exploratory analysis.....	21
8	Differences from protocol	21
9	References	21
9.1	Trial documents.....	21
9.2	Other references	22
10	Approval.....	23
11	Document history	24

Abbreviations

ADaM	Analysis Data Model
AE	Adverse event
CDISC	The Clinical Data Interchange Standards Consortium
CI	Confidence interval
COVID	Coronavirus-induced disease
CPAP	Continuous Positive Airway Pressure
CRP	C-reactive protein
DMC	Data Monitoring Committee
ECMO	Extra Corporeal Membrane Oxygenation
eCRF	Electronic case report form
ICD	International Classification of Diseases
ICNARC	Intensive Care National Audit and Research Centre
ITT	Intention to treat
MedDRA	Medical Dictionary for Regulatory Activities
OPCS-4	National Health Service OPCS Classification of Interventions and Procedures version 4
SARS	Severe acute respiratory syndrome
SARS-CoV-2	Severe acute respiratory syndrome coronavirus 2
SSAR	Suspected serious adverse reaction
SUSAR	Suspected unexpected serious adverse reaction
TSC	Trial Steering Committee

List of authors and reviewers (up to and including SAP version 1.1)

Authors

Dr Louise Linsell, Lead Trial Statistician, Nuffield Department of Population Health (NDPH), University of Oxford

Jennifer Bell, Trial Statistician, NDPH, University of Oxford

Reviewers

Professor Jonathan Emberson, Data Monitoring Committee (DMC) Statistician, NDPH, University of Oxford (prior to unblinded interim analysis of trial outcomes)

Professor Richard Haynes, Clinical Coordinator, NDPH, University of Oxford

Professor Peter Horby, Chief Investigator (CI), Nuffield Department of Medicine, University of Oxford

Professor Thomas Jaki, TSC Member, Department of Mathematics and Statistics, Lancaster University

Associate Professor Edmund Juszczak, TSC Member, NDPH, University of Oxford (until 6 July 2020)

Professor Martin Landray, Deputy CI, NDPH, University of Oxford

Professor Alan Montgomery, TSC Member, Nottingham Clinical Trials Unit, University of Nottingham

Dr Natalie Staplin, DMC Statistician, NDPH, University of Oxford (prior to unblinded interim analysis of trial outcomes)

List of authors and reviewers (version 2.0 onwards)

Authors

Professor Edmund Juszczak, TSC Member (University of Nottingham from 6 July 2020)

Professor Alan Montgomery (University of Nottingham), TSC Member

Professor Thomas Jaki (University of Cambridge) co-investigator and TSC Member

Reviewers

Enti Spata, Jennifer Bell, Dr Louise Linsell, Trial Statisticians, NDPH, University of Oxford

Professor Richard Haynes, Clinical Coordinator, NDPH, University of Oxford

Professor Martin Landray, Deputy CI, NDPH, University of Oxford

Professor Peter Horby, CI, Nuffield Department of Medicine, University of Oxford

Roles and responsibilities

Trial Statisticians

Until 30th September 2020: Dr Louise Linsell and Jennifer Bell (NDPH, University of Oxford)

Role: To develop the statistical analysis plan (blinded to trial allocation) and conduct the final comparative analyses for Lopinavir-Ritonavir, Corticosteroid (dexamethasone) and Hydroxychloroquine (main randomisation part A).

From 1st October 2020: Enti Spata (NDPH, University of Oxford)

Role: To develop the statistical analysis plan (blinded to trial allocation) and conduct the final comparative analyses for all other treatment arms.

Data Monitoring Committee (DMC) Statisticians

Professor Jonathan Emberson and Dr Natalie Staplin (NDPH, University of Oxford)

Role: To conduct regular interim analyses for the DMC. Contribution restricted up until unblinded to trial allocation.

Statisticians on the Trial Steering Committee (TSC)

Professor Edmund Juszcak (University of Nottingham), Professor Alan Montgomery (University of Nottingham), and Professor Thomas Jaki (University of Cambridge)

Role: Major organisational and policy decisions, and scientific advice; blinded to treatment allocation.

Trial IT systems & Programmers

Andy King, David Murray, Richard Welsh (NDPH, University of Oxford)

Role: To generate and prepare reports monitoring the randomisation schedule. To supply data snapshots for interim and final analysis. Responsibility for randomisation system, clinical databases and related activities.

Bob Goodenough (NDPH, University of Oxford)

Role: Validation of IT systems

Dr Will Stevens, Karl Wallendszuz (NDPH, University of Oxford)

Role: To produce analysis-ready datasets according to CDISC standards.

1 INTRODUCTION

This document details the proposed presentation and analysis for the main paper(s) reporting results from the multicentre randomised controlled trial RECOVERY (ISRCTN50189673) to investigate multiple treatments on major outcomes in inpatients for COVID-19 (clinically suspected or laboratory confirmed).

The results reported in these papers will follow the strategy set out here, which adheres to the guidelines for the content of a statistical analysis plan (SAP).¹ Any subsequent analyses of a more exploratory nature will not be bound by this strategy.

Suggestions for subsequent analyses by oversight committees, journal editors or referees, will be considered carefully in line with the principles of this analysis plan.

Any deviations from the statistical analysis plan will be described and justified in the final report. The analysis will be carried out by identified, appropriately qualified and experienced statisticians, who will ensure the integrity of the data during their processing.

This SAP is based on multiple versions of the protocol. All regulatory documents can be found in the RECOVERY trial directory: <https://www.recoverytrial.net/for-site-staff/site-set-up-1/regulatory-documents>.

SAP versions 1.0 & 1.1 applied to the first three principal comparisons (hydroxychloroquine, dexamethasone, and lopinavir-ritonavir versus no additional treatment respectively), for which data matured in the first UK wave of the pandemic. However, due to its later introduction, enrolment of patients in the azithromycin arm was much slower. Over time, factorial randomisations and a second randomisation have been added, introducing new treatment arms including convalescent plasma, tocilizumab, synthetic neutralizing antibodies, and aspirin. These changes, combined with the fact that use of corticosteroids (one of the original treatment arms) is now the usual standard of care for many patients, makes this a sensible juncture to re-evaluate the SAP and produce version 2.0.

2 BACKGROUND INFORMATION

2.1 Rationale

In early 2020, as the protocol was being developed, there were no approved treatments for COVID-19. The aim of the trial is to provide reliable evidence on the efficacy of candidate therapies (including re-purposed and novel drugs) for suspected or confirmed COVID-19 infection on major outcomes in hospitalised adult patients receiving standard care.

2.2 Objectives of the trial

2.2.1 *Primary objective*

To provide reliable estimates of the effect of study treatments on all-cause mortality within 28 days of the relevant randomisation.

2.2.2 *Secondary objectives*

To investigate the effect of study treatments on the duration of hospital stay and on the combined endpoint of use of invasive mechanical ventilation (including Extra Corporal Membrane Oxygenation [ECMO]) or death.

2.3 Trial design

This is a multi-centre, multi-arm, adaptive, open label, randomised controlled trial with three possible stages of randomisation, as described below. The trial is designed with streamlined processes in order to facilitate rapid large-scale recruitment with minimal data collection.

2.4 Eligibility

2.4.1 *Inclusion criteria*

Patients are eligible for the trial if all of the following are true:

- Hospitalised
- SARS-Cov-2 infection (clinically suspected or laboratory confirmed)
- No medical history that might, in the opinion of the attending clinician, put the patient at significant risk if they were to participate in the trial.

2.4.2 *Exclusion criteria*

If one or more of the active drug treatments is not available at the hospital or is believed, by the attending clinician, to be contraindicated (or definitely indicated) for the specific patient, then this fact will be recorded via the web-based form prior to randomisation; random allocation will then be between the remaining arms.

2.5 Treatments

All patients will receive standard management for the participating hospital. The main randomisation will be between the following treatment arms (although not all arms may be available at any one time). The doses listed are for adults; paediatric dosing is described in the protocol.

2.5.1 *Main randomisation part A:*

- **No additional treatment**
- **Lopinavir 400mg-Ritonavir 100mg** by mouth (or nasogastric tube) every 12 hours for 10 days. [Introduced in protocol version 1.0; **enrolment closed** 29 June 2020]

- **Corticosteroid** in the form of dexamethasone, administered as an oral liquid or intravenous preparation 6 mg once daily for 10 days. In pregnancy, prednisolone 40 mg administered by mouth (or intravenous hydrocortisone 80 mg twice daily) should be used instead. [Introduced in protocol version 1.0; **enrolment closed to adults 8 June 2020**]
- **Hydroxychloroquine** by mouth for 10 days (4 doses in first 24 hours and 1 dose every 12 hours for 9 days). [Introduced in protocol version 2.0; **enrolment closed 5 June 2020**]
- **Azithromycin 500mg** by mouth (or nasogastric tube) or intravenously once daily for a total of 10 days. [Introduced in protocol version 3.0; enrolment closed 27 November 2020]
- **Colchicine** by mouth for 10 days (1.5 mg in first 12 hours then 0.5 mg twice daily)

2.5.2 *Main randomisation part B:*

In a factorial design, eligible patients may be randomised to the arms below. The doses listed are for adults; paediatric dosing is described in the protocol.

- **No additional treatment**
- **Convalescent plasma** Single unit of ABO compatible convalescent plasma (275mls ± 75 mls) intravenous per day on study days 1 (as soon as possible after randomisation) and 2 (with a minimum of 12-hour interval between 1st and 2nd units). ABO identical plasma is preferred if available. The second transfusion should not be given if patient has a suspected serious adverse reaction during or after the first transfusion. [Introduced in protocol version 6.0; enrolment ongoing]
- **Synthetic neutralising antibodies** (REGN-COV2; adults and children aged ≥12 years only - children who weigh <40kg will also not be eligible for this treatment). A single dose of REGN10933 + REGN10987 8 g (4 g of each monoclonal antibody) in 250ml 0.9% saline infused intravenously over 60 minutes ± 15 minutes as soon as possible after randomisation. [Introduced in protocol version 9.1; enrolment ongoing]

2.5.3 *Main randomisation part C:*

In a factorial design, eligible patients may be randomised to the arms below. The dose listed is for adults; children are excluded from this comparison.

- **No additional treatment**
- **Aspirin** 150 mg by mouth (or nasogastric tube) or per rectum once daily until discharge. [Introduced in protocol version 10.1; enrolment ongoing]

2.5.4 *Second randomisation for patients with progressive COVID-19*

Patients enrolled in the main RECOVERY trial and with clinical evidence of a hyper-inflammatory state may be considered for a second randomisation if they meet the following criteria:

- Randomised into the main RECOVERY trial no more than 21 days ago
- Clinical evidence of progressive COVID-19:
 - oxygen saturation <92% on room air or requiring oxygen (or in children, significant systemic disease with persistent pyrexia, with or without evidence of respiratory involvement); and
 - C-reactive protein (CRP) ≥ 75 mg/L
- No medical history that might, in the opinion of the attending clinician, put the patient at significant risk if they were to participate in this aspect of the RECOVERY trial

Eligible participants may be randomised between the following treatment arms:

- **No additional treatment**
- **Tocilizumab** by intravenous infusion with the dose determined by body weight. [Introduced in protocol version 4.0; enrolment ongoing]

2.6 Definitions of primary and secondary outcomes

Outcomes will be assessed at 28 days and then 6 months after the relevant randomisation. Analysis of longer-term outcomes collected beyond this will be described in a separate Statistical Analysis Plan.

2.6.1 *Primary outcome*

Mortality (all-cause)

2.6.2 *Secondary clinical outcomes*

- Time to discharge from hospital
- Use of invasive mechanical ventilation (including Extra Corporal Membrane Oxygenation [ECMO]) or death (among patients not on invasive mechanical ventilation or ECMO at time of randomisation)

2.6.3 *Subsidiary clinical outcomes*

- Use of ventilation (overall and by type) among patients not on ventilation (of any type) at time of randomisation
- Duration of invasive mechanical ventilation among patients on invasive mechanical ventilation at time of randomisation (defined as time to successful cessation of invasive mechanical ventilation: see section 5.1.3.2)
- Use of renal dialysis or haemofiltration (among patients not on renal dialysis or haemofiltration at time of randomisation)
- Thrombotic events (overall and by type; introduced in Protocol version 10.1)

2.6.4 *Safety outcomes*

- Cause-specific mortality (COVID-19, other infection, cardiac, stroke, other vascular, cancer, other medical, external, unknown cause)

- Major cardiac arrhythmia (recorded on follow-up forms completed from 12 May 2020 onwards)
- Major bleeding (overall and by type; introduced in Protocol version 10.1)
- Early safety of antibody-based therapy (sudden worsening in respiratory status; severe allergic reaction; temperature $>39^{\circ}\text{C}$ or $\geq 2^{\circ}\text{C}$ rise since randomisation; sudden hypotension; clinical haemolysis; and thrombotic events within the first 72 hours; Main randomization phase B only)

2.6.5 *Detailed derivation of outcomes*

The detailed derivation of outcomes included in statistical analysis will be described separately in a data derivation document and included in the Study Data Reviewer's Guide.

2.7 Hypothesis framework

For each of the primary, secondary and subsidiary outcomes, the null hypothesis will be that there is no true difference in effect between any of the treatment arms.

2.8 Sample size

The larger the number randomised, the more accurate the results will be, but the numbers that can be randomised will depend critically on how large the epidemic becomes. If substantial numbers are hospitalised in the participating centres then it may be possible to randomise several thousand with moderate disease and a few thousand with severe disease. Some indicative sample sizes and projected recruitment will be estimated using emerging data for several different scenarios. Sample size and recruitment will be monitored by the TSC throughout the trial.

2.9 Randomisation

Eligible patients will be randomised using a 24/7 secure central web-based randomisation system, developed and hosted within NDPH, University of Oxford. Users of the system will have no insight into the next allocation, given that simple randomisation is being used. If a patient is randomised inadvertently more than once during the same hospital admission, the first allocation will be used.

The implementation of the randomisation procedure will be monitored by the Senior Trials Programmer, and the TSC notified if an error in the randomisation process is identified.

2.9.1 *Main randomisation part A*

Simple randomisation will be used to allocate participants to one of the following treatment arms (in addition to usual care), which is subject to change:

- No additional treatment
- Lopinavir-Ritonavir [Introduced in protocol version 1.0; **enrolment closed** 29 June 2020]
- Corticosteroid [Introduced in protocol version 1.0; **enrolment closed to adults** 8 June 2020]
- Hydroxychloroquine [Introduced in protocol version 2.0; **enrolment closed** 5 June 2020]

- Azithromycin [Introduced in protocol version 3.0; **enrolment closed 27 November 2020**]
- Colchicine [Introduced in protocol version 11.1]

The randomisation programme will allocate patients in a ratio of 2:1 between the no additional treatment arm and each of the other arms that are not contra-indicated and are available. Hence if all 4 active treatment arms are available, then the randomisation will be in the ratio 2:1:1:1:1. If one or more of the active drug treatments is not available at the hospital or is believed, by the attending clinician, to be contraindicated (or definitely indicated) for the specific patient, then this fact will be recorded via the web-based form prior to randomisation; random allocation will then be between the remaining arms (in a 2:1:1:1, 2:1:1 or 2:1 ratio).

2.9.2 *Main randomisation part B*

In a factorial design, eligible patients will be randomised simultaneously using simple randomisation with allocation ratio 1:1:1 to one of the following arms, which is subject to change:

- No additional treatment
- Convalescent plasma [Introduced in protocol version 6.0; enrolment ongoing]
- Synthetic neutralising antibodies [Introduced in protocol version 9.1; enrolment ongoing]

If the active treatment is not available at the hospital, the patient does not consent to receive convalescent plasma, or is believed, by the attending clinician, to be contraindicated for the specific patient, then this fact will be recorded via the web-based form and the patient will be excluded from the relevant arm in Randomisation part B.

2.9.3 *Main randomisation part C*

In a factorial design, eligible patients will be randomised simultaneously using simple randomisation with allocation ratio 1:1 to one of the following arms, which is subject to change:

- No additional treatment
- Aspirin [Introduced in protocol version 10.1; enrolment ongoing]

Note: From protocol version 7.0 onwards, randomisation is permitted in part B of main randomisation without randomisation in part A. From protocol version 10.1 onwards, randomisation is permitted in any combination of parts A, B, and C.

2.9.4 *Second randomisation for patients with progressive COVID-19*

Eligible participants will be randomised using simple randomisation with an allocation ratio 1:1 between the following arms, which is subject to change:

- No additional treatment
- Tocilizumab [Introduced in protocol version 4.0; enrolment ongoing]

2.10 Blinding

This is an open-label study. However, while the study is in progress, access to tabular results of study outcomes by treatment allocation will not be available to the research team, CIs, trial statisticians, clinical teams, or members of the TSC (unless the DMC advises otherwise). The DMC and DMC statisticians will be unblinded.

2.11 Data collection schedule

Baseline and outcome information will be collected on trial-specific electronic case report forms (eCRFs) and entered into a web-based IT system by a member of the hospital or research staff. Follow-up information will be collected on all study participants, irrespective of whether they complete the scheduled course of allocated study treatment. Study staff will seek follow-up information through various means, including routine healthcare systems and registries.

All randomised participants will be followed up until death or 6 months post-randomisation to the main trial (whichever is sooner). NHS Digital and equivalent organisations in the devolved nations will supply data fields relevant to trial baseline and outcome measures to NDPH, University of Oxford on a regular basis, for participants enrolled into the trial. This will be combined with the trial-specific data collected via the web-based IT system and adjudicated internally.

Longer term (up to 10 years) follow-up will be sought through linkage to electronic healthcare records and medical databases including those held by NHS Digital, Public Health England and equivalent bodies, and to relevant research databases (e.g. UK Biobank, Genomics England).

2.12 Data monitoring

During the study all study data will be supplied in strict confidence to the independent DMC for independent assessment and evaluation. The DMC will request such analyses at a frequency relevant to the emerging data from this and other studies.

The DMC has been requested to determine if, in their view, the randomised comparisons in the study have provided evidence on mortality that is strong enough (with a range of uncertainty around the results that is narrow enough) to affect national and global treatment strategies. Hence, multiple reviews by the Data Monitoring Committee have no material impact on the final analysis. In such a circumstance, the DMC will inform the TSC who will make the results available to the public and amend the trial arms accordingly.

2.13 Trial reporting

The trial will be reported according to the principles of the CONSORT statements.^{2, 3, 4} The exact composition of the trial publication(s) depends on the size of the epidemic, the availability of drugs, and the findings from the various pairwise comparative analyses (with the no additional treatment arm) in the main trial.

3 ANALYSIS POPULATIONS

3.1 Population definitions

The intention to treat (ITT) population will be all participants randomised, irrespective of treatment received. This ITT population will be used for analysis of efficacy and safety data. For interim analyses, baseline data will be reported for all participants with data available and outcome data will be reported for all participants who have died, been discharged from hospital, or reached day 28 after the first randomisation.

4 DESCRIPTIVE ANALYSES

4.1 Participant throughput

The flow of participants through the trial will be summarised for each separate pairwise comparison using a CONSORT diagram. The flow diagram will show the contribution of participants from each of the paths (from each of the parts of the main randomisation and from the second randomisation), where applicable. The flow diagrams will describe the numbers of participants randomly allocated, who received allocation, withdrew consent, and included in the ITT analysis population. The flow diagrams for arms in the main randomisation will also report the number of participants who underwent the second randomisation.

4.2 Baseline comparability of randomised groups

The following characteristics will be described separately for patients randomised to each main comparison (for each separate pairwise comparison of active treatment with the no additional treatment arm), and separately for the first and second randomisation.

4.2.1 Main randomisation (parts A, B and C)

- Age at randomisation
- Sex
- Ethnicity
- Region (UK, South East Asia)
- Time since COVID-19 symptoms onset
- Time since hospitalisation
- Current respiratory support
- Comorbidities (diabetes, heart disease, chronic lung disease, tuberculosis, human immunodeficiency virus, severe liver disease, severe kidney impairment)
- SARS-Cov-2 test result
- If female, known to be pregnant
- Use of systemic corticosteroid (including those allocated to corticosteroid in part A)
- Use of other relevant treatments (e.g. remdesivir, antiplatelet treatment, anticoagulant treatment)
- For part B only, anti-SARS-CoV-2 antibody concentration
- For treatment comparisons introduced in protocol v9.1 onwards:
 - C-reactive protein
 - Estimated glomerular filtration rate (calculated using the CKD-EPI formula)
 - D-dimer

4.2.2 Second randomisation

In addition to the above:

- Current respiratory support
- Latest oxygen saturation measurement
- Latest C-reactive protein
- Latest ferritin
- Latest estimated glomerular filtration rate (calculated using the CKD-EPI formula)
- Allocation in main randomisation parts A, B, and C
- Interval between first and second randomisation

The number and percentage will be presented for binary and categorical variables. The mean and standard deviation or the median and the interquartile range will be presented for continuous variables.

4.3 Completeness of follow-up

All reasonable efforts will be taken to minimise loss to follow-up, which is expected to be minimal as data collection for primary and secondary outcomes using trial-specific eCRFs is combined with linkage to routine clinical data on study outcomes from NHS Digital, ICNARC, and similar organisations in the devolved nations.

The number and percentage of participants with follow-up information at day 28 and at 6 months after the relevant randomisation will be reported. Data will be shown for each of the following: all-cause mortality, hospital discharge status, ventilation status, and will be shown for each randomised group for the main and second randomisation separately.

4.4 Adherence to treatment

The number and proportion of patients who did not receive the treatment they were allocated to will be reported. If any other trial treatment options were known to be received, instead of or in addition to, the allocated treatment during the 28-day follow-up period after the first randomisation, these will be collected and reported. Details on the number of days (or doses) of treatment received will be reported for all trial treatments received where available.

5 COMPARATIVE ANALYSES

For all outcomes, the primary analysis will be performed on the intention to treat (ITT) population at 28 days after randomisation. An ITT analysis of all outcomes at 6 months post-randomisation will also be conducted.

Pairwise comparisons will be made between each treatment arm and the no additional treatment arm (reference group) in that particular randomisation (main randomisation part A, main randomisation part B, main randomisation part C, and second randomisation). Since not all treatments may be available or suitable for all patients, those in the no additional

treatment arm will only be included in a given comparison if, at the point of their randomisation, they *could* alternatively have been randomised to the active treatment of interest (i.e. the active treatment was available at the time and it was not contra-indicated). The same applies to treatment arms added at a later stage; they will only be compared to those patients recruited concurrently.

5.1 Main randomisation part A

5.1.1 *Primary outcome*

Mortality (all-cause) will be summarised with counts and percentages by randomised comparison group. A time-to-event analysis will be conducted using the log-rank test, with the p-value reported. Kaplan-Meier estimates for the time to event will also be plotted (with associated log-rank p-values). The log-rank 'observed minus expected' statistic (and its variance) will be used to calculate the one-step estimate of the event rate ratio and confidence interval for each treatment group versus the no additional treatment group.⁵ For the primary outcome, discharge alive before the relevant time period (28 days after randomisation) will be assumed as absence of the event (unless there is additional data confirming otherwise).

5.1.2 *Secondary outcomes*

5.1.2.1 *Time to discharge alive from hospital*

A time-to-event analysis will be used to compare each treatment group with the no additional treatment group using the log-rank test. As described for the primary outcome, the rate ratio and its confidence interval will be estimated from the log-rank observed minus expected statistic and its variance, and Kaplan-Meier curves will be drawn. Patients who die in hospital will be censored after 28 days after randomisation. This gives an unbiased estimate of the recovery rate and comparable estimates to the competing risks approach in the absence of other censoring (which is expected to be very minimal).⁶

5.1.2.2 *Use of invasive mechanical ventilation (including ECMO) or death*

Counts and percentages will be presented by randomised group and the risk ratio will be calculated for each pairwise comparison with the no additional treatment arm, with confidence intervals and p-values reported. The absolute risk difference will also be presented with confidence intervals. Each component of this composite outcome will also be summarised. Patients who were already on invasive mechanical ventilation or ECMO at randomisation will be excluded from these analyses.

5.1.3 *Subsidiary clinical outcomes*

5.1.3.1 *Use of ventilation (overall and by type)*

Counts and percentages will be presented by randomised group for patients who received any assisted ventilation, together with risk ratios and confidence intervals for each pairwise comparison with the no additional treatment arm. The number of patients receiving the two main types of ventilation will also be reported: non-invasive ventilation (including CPAP, other non-invasive ventilation or high-flow nasal oxygen), and invasive mechanical ventilation

(including ECMO). Patients who were already receiving ventilation^a at randomisation will be excluded from these analyses.

5.1.3.2 Duration of invasive mechanical ventilation (time to successful cessation of invasive mechanical ventilation)

Successful cessation of invasive mechanical ventilation will be defined as removal of invasive mechanical ventilation within (and survival to) 28 days after randomisation. A time-to-event analysis will be used to compare each treatment group with the no additional treatment group using the log-rank test, as described above. The rate ratio and its confidence interval will be estimated from the log-rank observed minus expected statistic and its variance, and Kaplan-Meier curves will be drawn. Patients who die within 28 days of randomisation will be censored *after* 28 days after randomisation. Patients who were not already on invasive mechanical ventilation or ECMO at randomisation will be excluded from these analyses.

5.1.3.3 Use of renal dialysis or haemofiltration

Counts and percentages will be presented by randomised group and the risk ratio will be calculated for each pairwise comparison with the no additional treatment arm, with confidence intervals and p-values reported. The absolute risk difference will also be presented with confidence intervals. Patients who were already on renal dialysis or haemofiltration at randomisation will be excluded from these analyses.

5.1.3.4 Thrombotic event

Counts and percentages will be presented by randomised group. The absolute risk differences will also be presented with confidence intervals. Type of thrombotic event will also be described: (i) acute pulmonary embolism; (ii) deep vein thrombosis; (iii) ischaemic stroke, (iv) myocardial infarction; (v) systemic arterial embolism; and (vi) all sites combined.

5.2 Main randomisation part B

In the factorial design, the main effects of treatments evaluated in part B will be presented and tested across all arms in main randomisation parts A and C combined, as described in 5.1. (Assessments of whether the effects of treatments in part B vary depending on other randomised treatments are described in section 5.7).

5.3 Main randomisation part C

In the factorial design, the main effects of treatments evaluated in part C will be presented and tested across all arms in main randomisation parts A and B combined, as described in 5.1. (Assessments of whether the effects of treatments in part C vary depending on other randomised treatments are described in section 5.7).

5.4 Second randomisation

^a For comparisons introduced to the main randomisation prior to protocol version 9.1, patients who were already receiving oxygen at randomisation will also be excluded from these analyses (since it is not possible to distinguish those who were already receiving non-invasive ventilation).

Evaluation of treatment effects in the main randomisation and the second randomisation will be conducted independently, as described in 5.1.

5.5 Pre-specified subgroup analyses

Pre-specified subgroup analyses will be conducted for the main randomisation (parts A, B and C) and the second randomisation, for the following outcomes:

- Mortality (all-cause)
- Time to discharge from hospital
- Use of invasive mechanical ventilation (including ECMO) or death

Tests for heterogeneity (or tests for trend for 3 or more ordered groups) will be conducted to assess whether there is any good evidence that the effects in particular subgroups differ materially from the overall effect seen in all patients combined. Results will be presented on forest plots as event rate ratios, or risk ratios, with confidence intervals. The following subgroups will be examined based on information at randomisation:

- Age (<70; 70-79; 80+ years)
- Sex (Male; Female)
- Ethnicity (White; Black, Asian or Minority Ethnic)
- Region (UK, South East Asia)
- Time since illness onset (≤ 7 days; > 7 days)
- Requirement for respiratory support
 - For main randomisation: None; Oxygen only (with or without non-invasive ventilation); Invasive mechanical ventilation (including ECMO)
 - For second randomisation: No ventilator support (including no or low-flow oxygen); Non-invasive ventilation (including CPAP, other non-invasive ventilation, or high-flow nasal oxygen), Invasive mechanical ventilation (including ECMO)
- Use of systemic corticosteroid (including dexamethasone)
- For part B only: Recipient anti-SARS-CoV-2 antibody concentration at randomisation ($< 8 \times 10^6$ units; $\geq 8 \times 10^6$ units^b)

5.6 Sensitivity analyses

Sensitivity analyses of the primary and secondary outcomes will be conducted among those patients with a positive test for SARS-COV-2 (i.e. confirmed cases).

5.7 Other exploratory analyses

In addition, exploratory analyses will be conducted to test for interactions between treatments allocated in each of the different randomisations, provided that doing so does not lead to premature unblinding of results for ongoing comparators.

^b Measured by Oxford immunoassay. 8×10^6 represents the threshold shown to distinguish evidence of prior infection from no evidence of prior infection with sensitivity and specificity $> 98\%$.⁷ Exploratory analyses using other assays may determine appropriate cut-offs based on blinded inspection of associations between assay results and baseline characteristics and mortality.

Non-randomised exploratory analyses will be used to explore the likely influence of different levels of convalescent plasma antibody concentration on the efficacy of convalescent plasma.

Additional analyses will set the results for children (<18 years) and pregnant women in the context of the overall results.

5.8 Adjustment for baseline characteristics

The main analyses described above will be unadjusted for baseline characteristics. However, if there are any important imbalances between the randomised groups in key baseline pre-specified subgroups (see section 5.4) or allocation in the orthogonal components of the main randomisation, where applicable, emphasis will be placed on analyses that are adjusted for the relevant baseline characteristic(s). This will be done using Cox regression for the estimation of adjusted hazard ratios and a log-binomial regression model for the estimation of adjusted risk ratios.

5.9 Significance levels and adjustment of p-values for multiplicity

Evaluation of the primary trial (main randomisation) and secondary randomisation will be conducted independently, and no adjustment be made for these. Formal adjustment will not be made for multiple treatment comparisons, the testing of secondary and subsidiary outcomes, or subgroup analyses. However, due allowance for multiple testing will be made in the interpretation of the results: the larger the number of events on which a comparison is based and the more extreme the P-value after any allowance has been made for the nature of the particular comparison (i.e. primary or secondary; pre-specified or exploratory), the more reliable the comparison and, hence, the more definite any finding will be considered. 95% confidence intervals will be presented for the main comparisons.

5.10 Statistical software employed

The statistical software SAS version 9.4 and R Studio 3.6.2 (or later) for Windows will be used for the interim and final analyses.

5.11 Data standards and coding terminology

Datasets for analysis will be prepared using CDISC standards for SDTM and ADaM. Wherever possible, clinical outcomes (which may be obtained in a variety of standards, including ICD10 and OPCS-4) will be coded using MedDRA version 20.1.

6 SAFETY DATA

Suspected serious adverse reactions (SSARs) and suspected unexpected serious adverse reactions (SUSARs) will be listed by trial allocation.

For each of the following, counts and percentages will be presented by randomised group. Where possible, the absolute risk differences will also be presented with confidence intervals:

6.1.1.1 *Cause-specific mortality*

Cause-specific mortality (COVID-19, other infection, cardiac, stroke, other vascular, cancer, other medical, external, unknown cause) will be analysed in a similar manner to the primary outcome.

6.1.1.2 *Major cardiac arrhythmia*

Type of arrhythmia will also be described: (i) atrial flutter or fibrillation; (ii) supraventricular tachycardia; (iii) ventricular tachycardia; (iv) ventricular fibrillation; (v) atrioventricular block requiring intervention, with subtotals for (i)-(ii) and (iii)-(iv).

6.1.1.3 *Major bleeding*

Type of bleeding will also be described: (i) intracranial bleeding; (ii) gastro-intestinal bleeding; (iii) other bleeding site, and (iv) all sites combined.

6.1.1.4 *Early safety of antibody-based therapy*

Additional safety data will be collected in a subset of patients randomised to part B: (i) sudden worsening in respiratory status; (ii) severe allergic reaction; (iii) temperature $>39^{\circ}\text{C}$ or $\geq 2^{\circ}\text{C}$ rise since randomisation; (iv) sudden hypotension; (v) clinical haemolysis; and (vi) thrombotic event.

7 ADDITIONAL POST-HOC EXPLORATORY ANALYSIS

Any post-hoc analysis requested by the oversight committees, a journal editor or referees will be labelled explicitly as such. Any further future analyses not specified in the analysis protocol will be exploratory in nature and will be documented in a separate statistical analysis plan.

8 DIFFERENCES FROM PROTOCOL

The testing of multiple treatment arms will not formally be adjusted for, but given the number of comparisons, due allowance will be made in their interpretation. Formal methods of adjustment for multiplicity were not adopted because of treatment arms being added over time (including the factorial convalescent plasma comparison), unequal recruitment into each arm, and the ultimate number of treatments under evaluation not known in advance.

This analysis plan will be updated prior to unblinding of the 6-month follow-up results. Additional analyses may be specified, e.g. to explore the impact of randomised treatment allocation on hospital re-admission for COVID-19.

9 REFERENCES

9.1 Trial documents

Study protocol, case report forms, training materials, and statistical analysis plan are published on the trial website.

9.2 Other references

1. Gamble C, Krishan A, Stocken D, Lewis S, Juszczak E, Doré C, Williamson PR, Altman DG, Montgomery A, Lim P, Berlin J, Senn S, Day S, Barbachano Y, Loder E. Guidelines for the Content of Statistical Analysis Plans in Clinical Trials. *JAMA* 2017;318(23):2337-2343.
2. Schulz KF, Altman DG, Moher D for the CONSORT Group. CONSORT 2010 Statement: updated guidelines for reporting parallel group randomised trials. *BMJ* 2010;340:698-702.
3. Juszczak E, Altman DG, Hopewell S, Schulz KF. Reporting of multi-arm parallel-group randomized trials: extension of the CONSORT 2010 statement. *JAMA* 2019;321(16):1610-1620.
4. Dimairo M, Pallmann P, Wason J, Todd S, Jaki T, Julious SA, Mander AP, Weir CJ, Koenig F, Walton MK, Nicholl JP, Coates E, Biggs K, Hamasaki T, Proschan MA, Scott JA, Ando Y, Hind D, Altman DG; ACE Consensus Group. The Adaptive designs CONSORT Extension (ACE) statement: a checklist with explanation and elaboration guideline for reporting randomised trials that use an adaptive design. *BMJ*. 2020 Jun 17;369:m115. doi: 10.1136/bmj.m115. PMID: 32554564; PMCID: PMC7298567.
5. Peto R, Pike MC, Armitage P, Breslow NE, Cox DR, Howard SV, Mantel N, McPherson K, Peto J, Smith PG. Design and analysis of randomized clinical trials requiring prolonged observation of each patient. Part II: analysis and examples. *Br J Cancer* 1977;35:1-39.
6. Betensky RA and Schoenfeld DA. Nonparametric Estimation in a Cure Model with Random Cure Times. *Biometrics* 2001;57:282-286.
7. The National SARS-CoV-2 Serology Assay Evaluation Group. Performance characteristics of five immunoassays for SARS-CoV-2: a head-to-head benchmark comparison. *Lancet Infect Dis* 2020; 20: 1390-1400

10 APPROVAL

Trial Statistician	Name: Mr Enti Spata	
	Signature:	Date:
Chief Investigator	Name: Professor Peter Horby	
	Signature:	Date:
Deputy Chief Investigator	Name: Professor Martin Landray	
	Signature:	Date:
Steering Committee Statistician	Name: Professor Edmund Juszcak	
	Signature:	Date:
Steering Committee Statistician	Name: Professor Alan Montgomery	
	Signature:	Date:
Steering Committee Statistician	Name: Professor Thomas Jaki	
	Signature:	Date:

11 DOCUMENT HISTORY

Version	Date	Edited by	Comments/Justification	Timing in relation to unblinded interim monitoring	Timing in relation to unblinding of Trial Statisticians
0.1	20/03/20	LL/JB	First draft.	Prior	Prior
0.2	01/04/20	LL/JB	Comments and amendments from Martin Landray, Jonathan Emberson & Natalie Staplin. Also aligned with updated protocol and CRFs.	Prior	Prior
0.3	01/04/20	EJ/LL	Further edits and comments.	Prior	Prior
0.4	07/04/20	JB/EJ/LL	Following statistics group meeting on 02/04/20.	Prior	Prior
0.5	22/04/20	JB/LL/EJ	Following statistics group meeting on 09/04/20 and further protocol update.	After	Prior
0.6	24/04/20	LL	Following statistics group meeting on 23/04/20.	After	Prior
0.7	10/05/20	LL	Protocol update.	After	Prior
0.8	15/05/20	LL	Following statistics group meeting on 15/05/20.	After	Prior
0.9	27/05/20	LL	Further comments from TSC members prior to interim analysis on 28/05/20.	After	Prior
1.0	09/06/20	LL	Revised following the stopping of the hydroxychloroquine arm, and prior to the trial statisticians receiving unblinded data for this arm.	After	Prior
1.1	21/06/20	LL/JB/RH	Additional clarification of ventilation denominators. Adjustment for any imbalances of subgroup characteristics between treatment arms at randomisation. Clarification of analysis of composite outcome. Removal of 'Unknown' ethnicity subgroup. Addition of section 5.5 Adjustment for baseline characteristics.	After	After unblinding of hydroxychloroquine and dexamethasone arms.

Version	Date	Edited by	Comments/Justification	Timing in relation to unblinded interim monitoring	Timing in relation to unblinding of Trial Statisticians
2.0	04/11/20	EJ/ES	Revised to reflect changes in protocol, including introduction of factorial randomisations and new arms, including convalescent plasma, tocilizumab, synthetic neutralizing antibodies (REGN-COV2, and aspirin.	Prior to interim analysis of aspirin arm After interim analyses of all other arms	After unblinding of 28-day results for hydroxychloroquine, lopinavir-ritonavir, and dexamethasone arms. Prior to unblinding of any other arms
2.1	02/12/20	ES	Addition of colchicine. Modification of definition of recipient antibody concentration subgroup.	Prior to interim analyses including antibody results or of colchicine arm.	After unblinding of 28-day results for hydroxychloroquine, lopinavir-ritonavir, and dexamethasone arms. Prior to unblinding of any other arms